



A service of the National Library of Medicine
and the National Institutes of Health

My NCBI
[\[Sign In\]](#)

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals

Search PubMed for electromagnetic radio cancer Go Clear

Limits Preview/Index History Clipboard Details

About Entrez
NCBI Toolbar

Note: Performing your original search, *electromagnetic radio cancer*, in PubMed will retrieve [122 citations](#).

Text Version

Display Abstract Show 20 Sort by Send to

Entrez PubMed

Overview
Help | FAQ
Tutorials
New/Noteworthy
E-Utilities

All: 1 Review: 0

1: [Eur J Cancer Prev.](#) 2003 Aug;12(4):301-7.

[Related Articles, Links](#)



Follow-up of radio and telegraph operators with exposure to electromagnetic fields and risk of breast cancer.

[Kliukiene J](#), [Tynes T](#), [Andersen A](#).

The Cancer Registry of Norway, Institute of Population-based Cancer Research, Montebello, NO-0310 Oslo, Norway.
jolanta.kliukiene@krefregisteret.no

It is still unclear whether exposure to electromagnetic fields (EMF) is associated with breast cancer. To further investigate the issue, we followed-up a cohort of Norwegian female radio and telegraph operators in the period 1 January 1961 to 31 May 2002, with 99 breast cancer cases. The standardized incidence ratio (SIR) for breast cancer was 1.30 (95% confidence intervals (CI) 1.05-1.58), compared with the total Norwegian female population. In a subsequent nested case-control study, exposure to radio frequency (405 kHz-25 MHz) and extremely low-frequency (50 Hz) fields due to stay in the radio room during day and night was cumulated by years of employment and workload according to ship type. The exposure was assessed in two age groups (<50, 50+) with regard to risk of breast cancer. The odds ratios in the group with the highest cumulative exposure were 1.78 (95% CI 0.59-5.41) and 2.37 (95% CI 0.88-6.36) in the younger and the older women, respectively. P-value for trend was 0.03 in both age groups. The results of the oestrogen receptor status analysis by exposure to EMF showed an increased risk of oestrogen receptor-positive breast cancer in the younger women, while the older age group had an elevated risk of oestrogen receptor-negative breast cancer. Thus, the present study contributes to the hypothesis of an association between occupational exposure to EMF and increased risk of breast cancer.

PMID: 12883383 [PubMed - indexed for MEDLINE]

Display	Abstract	Show	20	Sort by	Send to
---------	----------	------	----	---------	---------

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
Department of Health & Human Services
[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Feb 13 2006 12:53:38